

Self-energy and vertex radiative corrections in LQG

Claudio Perini^{ab}, Carlo Rovelli^a, Simone Speziale^{ac}

^aCentre de Physique Théorique de Luminy*, Case 907, F-13288 Marseille, EU

^bDipartimento di Matematica, Università Roma Tre, Largo Murialdo 1, I-00146 Roma, EU

^cPerimeter Institute, 31 Caroline St North, Waterloo, ON N2L 2Y5, Canada

(Dated: October 10, 2008)

We consider the elementary radiative-correction terms in loop quantum gravity. These are a two-vertex “elementary bubble” and a five-vertex “ball”; they correspond to the one-loop self-energy and the one-loop vertex correction of ordinary quantum field theory. We compute their naive degree of (infrared) divergence.

I. INTRODUCTION

Recent developments in the spinfoam research program [1, 2, 3, 4, 5] have provided a novel and manageable perturbative definition of the dynamics of loop quantum gravity [6, 7]. The theory is first cut-off by choosing a 4d triangulation Δ_N of spacetime, formed by N 4-simplices; then the continuous theory can be defined by the $N \rightarrow \infty$ limit of the expectation values.¹ At fixed N , the partition function of the theory is given by a sum over spins and intertwiners, which can be interpreted as a version of the Misner-Hawking “sum over geometries” [9], regulated by replacing continuous geometries with Regge geometries [10] on Δ_N . This sum may contain divergent terms. Here we study these terms. For previous studies of divergences in spinfoam models see [11, 12, 13, 14].

We consider the version of the formalism defined in [5], which coincides with the one discussed in [3] when the Immirzi parameter $\gamma \in \mathbb{R}^+$ satisfies $\gamma < 1$. The theory is defined by the partition function

$$Z = \sum_{j_f, i_e} \prod_f d(j_f) \prod_v A_v(j_f, i_e). \quad (1)$$

The sum is over $SU(2)$ irreducible representations j_f associated to the triangles f of Δ_N and $SU(2)$ intertwiners i_e associated to tetrahedra e of Δ_N . $d(j)$ is a face amplitude and the vertex amplitude is given by

$$A_v(j_f, i_e) \equiv \lambda \sum_{i_e^+, i_e^-} 15j_N(j_f^+, i_e^+) 15j_N(j_f^-, i_e^-) \prod_e f_{i_e^+, i_e^-}^{i_e}, \quad (2)$$

where $j^\pm \equiv \frac{1 \pm \gamma}{2} j$; $15j_N$ denotes the contraction of five normalized 4-valent intertwiners according to the pattern of a 4-simplex (see (A19)), and $f_{i_e^+, i_e^-}^{i_e}$ are the fusion coefficients [1], discussed in detail below. The arguments of A_v are the ten spin and the five intertwiners (labeled by virtual spins in a coupling channel) adjacent to the vertex v . We have inserted a dimensionless coupling constant λ . This vertex amplitude is derived in [1, 2, 3, 4, 5] from the action of GR; see these references for details. On the semiclassical behavior of the sum (1), see [15].

Transition amplitudes can be constructed by picking a triangulation Δ_N with a boundary $\partial\Delta_N$, and fixing spins and edges on the boundary in (1):

$$W(j_l, i_n) = \sum_{j_f, i_e} \prod_f d(j_f) \prod_v A_v(j_f, i_e), \quad (3)$$

where j_l are the spins of the boundary faces (or “links”) and i_n are the intertwiners of the boundary tetrahedra (or “nodes”). (The expressions “links” and “nodes” refer to the graph Γ defined by the 1-skeleton of the cellular complex dual to $\partial\Delta_N$.) We are interested in the potential divergences of the sums (3).

As explained in the next Section, the structure of such divergences bear some similarities (but also some differences) with that of the divergences in Feynman diagrams. There, divergences come from momentum integrals associated to

* Unité mixte de recherche (UMR 6207) du CNRS et des Universités de Provence (Aix-Marseille I), de la Méditerranée (Aix-Marseille II) et du Sud (Toulon-Var); laboratoire affilié à la FRUMAM (FR 2291).

¹ This is similar to the well-known nonperturbative definition of QCD via a lattice [8], but with the difference that because of general covariance the lattice spacing does not affect physical expectation values.

internal *loops*. Here, divergences turn out to be associated to *bubbles* in the triangulation. A *bubble* is a collection of faces in the 2-skeleton of the cellular complex dual to Δ_N , forming a closed 2-surface. As a first step towards the characterization of all divergences, we study here the most elementary of such bubbles, and the potential divergences they give rise to. In particular, we focus on two elementary graphs: the self-energy (or “elementary bubble”), and the vertex divergence (or “elementary ball”). As in QFT, these can be viewed as divergences on particularly simple triangulations with boundaries (a Δ_2 and a Δ_5), or as divergences arising in sub-triangulations in larger triangulations.

It is shown in [16] that any spinfoam model can be expressed as a Feynman amplitude of a group field theory [17]. If we view (3) as determined by the Feynman expansion of a group field theory, these graphs correspond precisely to the self-energy correction to the propagator and the first radiative correction of the vertex, from which, in several QFT’s, all other divergences depend.

Our results are summarized below, in Section V. We find that the naive degree of divergence depends crucially on the details of the normalization of the fusion coefficients mentioned above, and the faces amplitude. It is therefore important to gain a better control over these quantities, which are not clearly fixed in the literature. For an appropriate and natural choice of the normalization of the fusion coefficients, both diagrams are finite.

II. PRELIMINARIES

A. Bubbles

At fixed spins j_f , the sum over the intertwiners i_e in (3) is always over a finite number of terms. Therefore divergences can only arise from $j_f \rightarrow \infty$. The spins j_f are constrained by the Clebsch-Gordan inequalities at each edge, because when these relations are not satisfied the vertex amplitude vanishes. This fact determines the structure of the possible divergences [11, 12].

A well-known way of resolving the Clebsch-Gordan inequalities, indeed, is to change variables as follows. Replace a face with spin j_f by $2j_f$ overlapping sheets. At a given edge e , the Clebsch-Gordan conditions are satisfied if and only if there is a way of pairing all individual sheets coming from the four faces that join at the edge, in such a way that each sheet is paired with one and only one sheet of a different face. The pairing joins individual sheets across faces, forming surfaces that wrap over the 2-skeleton of the triangulation. Such surfaces can be of two kinds: either they meet the boundary, or they are closed. Call the closed surfaces “bubbles”. Suppose we want to increase the spins j_f to infinity (keeping the boundary spins j_l fixed). Then we have to add more and more of such surfaces. However, we cannot add surfaces that meet the boundary, because this would increase j_l . Therefore the only possibility is to increase the number of surfaces of the closed kind, or “bubbles” [11, 12].

The situation is thus similar to that in conventional QFT, where momentum conservation at QFT vertices implies that divergences are necessarily associated with closed lines on the Feynman graph, or “loops”. Here Clebsch-Gordan inequalities at each edge imply that divergences are necessarily associated with “bubbles”.

B. Normalizations

The convergence properties of (1) depend on both face and vertex amplitudes. The recent developments in the spinfoam dynamics have resulted in a vertex amplitude that, at least for $\gamma < 1$, is unique up to normalization; while the face amplitudes are not fully fixed yet. At the present stage of investigation, it is therefore useful to study the convergence properties of a family of models like (1), which differ by the face amplitudes and the normalization of the vertex.

The face amplitude $d(j)$ in (1) was taken in [5] to be the dimension of the $SO(4)$ representation determined (via the Plebanski constraints) by the $SU(2)$ representation j

$$d(j) = d_{SO(4)}(j) \equiv (2j^- + 1)(2j^+ + 1) = (|1 - \gamma|j + 1)((1 + \gamma)j + 1) \quad (4)$$

on the basis of an analogy with BF theory. However, this part of the definition of the theory is the most uncertain. A natural alternative is to chose the dimension of the representation j itself

$$d(j) = d_{SU(2)}(j) \equiv (2j + 1). \quad (5)$$

To keep the issue open, we consider here a generic dependence such that

$$d(j) \sim j^k \quad (6)$$

when $j \rightarrow \infty$. The two cases considered above correspond to $k = 2$ and $k = 1$, respectively.

The fusion coefficients entering (2) are covariant maps from the $SO(4)$ intertwiner space to the $SO(4)$ intertwiner space. They were defined in [1] as

$$f_{i^+i^-}^i(j_a) = \langle i^+i^- | f | i \rangle = i^{abcd} C_a^{a^+a^-} C_b^{b^+b^-} C_c^{c^+c^-} C_d^{d^+d^-} i_{a^+b^+c^+d^+}^+ i_{a^-b^-c^-d^-}^-. \quad (7)$$

where i^{abcd} are normalized 4-valent intertwiners and $C_a^{a^+a^-} = \langle j_1^+ j_1^-, a^+ a^- | j_1 a \rangle$ are Clebsch-Gordan coefficients. See Appendix A for precise definitions. (The group integral appearing in the definition in [1] is not needed once we project on the $SO(4)$ invariant subspace spanned by the $|i^+i^- \rangle$ intertwiners.) It is not clear to us what are the conditions that fix the normalization of the fusion coefficients defined in [1]. In particular, there is a more natural alternative given by

$$\tilde{f}_{i^+i^-}^i(j_a) = \langle i^+i^- | f | i \rangle = i^{abcd} i_a^{a^+a^-} i_b^{b^+b^-} i_c^{c^+c^-} i_d^{d^+d^-} i_{a^+b^+c^+d^+}^+ i_{a^-b^-c^-d^-}^- \quad (8)$$

where i_a^{bc} are normalized trivalent intertwiners (see Appendix); these fusion coefficients are related to the above ones by

$$f_{i^+i^-}^i(j_a) = \sqrt{\prod_{a=1\dots 4} (2j_a + 1)} \tilde{f}_{i^+i^-}^i(j_a). \quad (9)$$

This uncertainty in the normalizations is similar to the one in old spinfoam models: see [11, 12, 13, 14] and [6].

III. THE ELEMENTARY BUBBLE

Consider a triangulation Δ_2 formed by two 4-simplices joined along four of their boundary tetrahedra. Its dual complex is illustrated in Figure 1. The boundary triangulation $\partial\Delta_2$ is formed by two tetrahedra, joined by all their triangles. The boundary graph is therefore formed by two nodes joined by four links. Denote i_u, i_d the intertwiners on the nodes and j_a , where $a = 1, \dots, 4$, the spins of the four links (see Figure 1). These are the boundary variables. The internal variables are the four intertwiners i_a on the four internal tetrahedra, and the six spins j_{ab} on the six internal triangles (see Figure 2). The amplitude is

$$W(j_a, i_u, i_d) = \lambda^2 \sum_{j_{ab}, i_a} \prod_a d(j_a) \prod_{ab} d(j_{ab}) A_v(j_a, j_{ab}, i_a, i_u) A_v(j_a, j_{ab}, i_a, i_d). \quad (10)$$

We take $\lambda = 1$ for simplicity in what follows. To study if this amplitude diverges, it is convenient to set all external spins j_a to zero. (This is analogous to the analysis of the vertex divergences in, say, $\lambda\phi^4$ QFT performed by setting the four external momenta to zero.) In this case, $i_u = i_d = 0$, where “0” indicates the trivial intertwiners between trivial representations, and we have

$$W_2 \equiv W(0, 0, 0) = \sum_{j_{ab}, i_a} \prod_{ab} d(j_{ab}) A_v^2(0, j_{ab}, i_a, 0). \quad (11)$$




FIG. 1: The self-energy spinfoam. Left: boundary variables. Right: internal variables (only one of the six internal faces is highlighted).




FIG. 2: The self-energy bubble. Left: the six faces. Right: equatorial section.

A. Simple example: $SU(2)$ BF

Before performing the computation for the gravity case, let us start with the simpler calculation for a (topological) four-dimensional $SU(2)$ BF theory. In this case,

$$d(j) = (2j + 1)^k \quad (12)$$

where the topological theory is obtained for $k = 1$; and

$$A_v(j_{nn'}, i_n) = 15j_N(j_{nn'}, i_n). \quad (13)$$

When we set four spins sharing the same node to zero the intertwiners are all trivial, and the normalized $15j_N$ reduces to the $6j$ symbol for the remaining six spins (see the Appendix.) Thus (11) becomes

$$W_2 = \sum_{j_{ab}} \prod_{ab} (2j_{ab} + 1)^k (6j(j_{ab}))^2 \quad (14)$$

To estimate the degree of divergence of the above quantity we look at the large spin behaviour of the summand. Using the Ponzano-Regge asymptotic expression for the $6j$ symbol [18]

$$6j \sim \frac{1}{\sqrt{V}} (e^{iS} + e^{-iS}), \quad (15)$$

we have

$$(6j)^2 \sim \frac{1}{V} (2 + e^{2iS} + e^{-2iS}) \sim \frac{1}{V}. \quad (16)$$

Here V is the volume of a tetrahedron that has the j_{ab} as lengths. If $j_{ab} \sim j \rightarrow \infty$, then $V \sim j^3$ and $(6j)^2 \sim j^{-3}$. Combining with the measure factor gives

$$W_2 \sim \sum_{j_{ab}} j^{6k-3}. \quad (17)$$

The naive degree of divergence is given by the power of the summand plus the number of sums. The Clebsch-Gordan conditions do not change this, because the volume of the region of $j_{ab} < \Lambda$ satisfying these relations still grows like Λ^6 . Hence

$$D_{\text{BF}} = 6 + 6k - 3. \quad (18)$$

The graph converges only if $k < -\frac{1}{2}$. For the conventional $k = 1$ it diverges as Λ^9 , with Λ an infrared cut off.

For $k = 1$, an alternative way to obtain the same estimate is the following. The expression (14) is precisely the Ponzano-Regge partition function of a triangulation formed by two tetrahedra glued by all their faces, and it is equivalent to

$$W_2 = \int dU_a \prod_{ab} \delta(U_a U_b^{-1}). \quad (19)$$

Integrating out the delta functions explicitly one by one, leaves

$$W_2 = (\delta(\mathbb{1}))^3. \quad (20)$$

Using the Plancherel expansion of the δ function (see (A13) in the Appendix) this becomes

$$W_2 = \left(\sum_j (2j + 1) \text{tr}[R^{(j)}(\mathbb{1})] \right)^3 = \left(\sum_j (2j + 1)^2 \right)^3 \quad (21)$$

which indeed diverges as Λ^9 , confirming the result above. We now return to gravity.

B. Gravity

Inserting the gravitational vertex amplitude (2) into equation (10) gives

$$W_2 = \sum_{j_{ab}, i_a} \prod_{ab} d(j_{ab}) \left(\sum_{i_n^+, i_n^-} 15j_N(0, \frac{1+\gamma}{2}j_{ab}, i_n^+) 15j_N(0, \frac{1-\gamma}{2}j_{ab}, i_n^-) \prod_n f_{i_n^+, i_n^-}^{i_n} \right)^2. \quad (22)$$

Choosing the virtual spin i in the pairing $((j_1, j_2), (j_3, j_4))$ to label the intertwiners, fusion coefficients can be written in the form [20]

$$f_{i^+ i^-}^i(j_1, j_2, j_3, j_4) = \left[(2i+1)(2i^++1)(2i^-+1) \prod_{n=1}^4 (2j_n+1) \right]^{\frac{1}{2}} \left\{ \begin{matrix} j_1^+ & i^+ & j_2^+ \\ j_1^- & i^- & j_2^- \\ j_1 & i & j_2 \end{matrix} \right\} \left\{ \begin{matrix} j_3^+ & i^+ & j_4^+ \\ j_3^- & i^- & j_4^- \\ j_3 & i & j_4 \end{matrix} \right\}, \quad (23)$$

where the matrices are Wigner $9j$ -symbols (other choices of pairings are equivalent as W_2 does not depend on the basis chosen.) As above, the presence of four vanishing spins fixes the intertwiners and reduces the $15j_N$ -symbol to the Wigner $6j$ -symbol. One of the fusion coefficient (the one among all trivial representations) is equal to unity. We have then

$$W_2 = \sum_{j_{ab}, i_a} \prod_{ab} d(j_{ab}) \left(6j(\frac{1+\gamma}{2}j_{ab}) \ 6j(\frac{1-\gamma}{2}j_{ab}) \prod_a f_a \right)^2 \quad (24)$$

where f_a stands for

$$f_1 = \sqrt{d(j_{12})d(j_{13})d(j_{14})} \left\{ \begin{matrix} j_{12}^+ & j_{14}^+ & j_{13}^+ \\ j_{12}^- & j_{14}^- & j_{13}^- \\ j_{12} & j_{14} & j_{13} \end{matrix} \right\} \quad (25)$$

and so on cyclically. We show in Appendix B that for large $j_{ab} \simeq j$, the f 's behave like $j^{-3/4}$. Together with the scaling (16), we get

$$W_2 \sim \sum_{j_{ab}} (j^k)^6 [(j^{-3/2})^2 (j^{-3/4})^4]^2. \quad (26)$$

We see that the degree of divergence with the fusion coefficients f in (7) is

$$D_{W_2} = 6 + 6k - 12. \quad (27)$$

Naively, the self-energy diverges as a power for $k = 2$, and logarithmically for $k = 1$.

Next, we consider the fusion coefficients normalized as in (8). As shown in the Appendix, these scale like $j^{-9/4}$. We then get

$$\widetilde{W}_2 \sim \sum_{j_{ab}} (j^k)^6 [(j^{-3/2})^2 (j^{-9/4})^4]^2. \quad (28)$$

Therefore the degree of divergence with the fusion coefficients $\tilde{f}$ is

$$D_{\widetilde{W}_2} = 6 + 6k - 24, \quad (29)$$

which converges for $k < 3$. The stronger convergence could have been anticipated from (9).




FIG. 3: The “ball” vertex amplitude. Left: the five vertices and the ten+five edges. Right: one internal face and one external face are shown.

IV. THE BALL

The next interesting diagram is a 4-simplex expanded with a 1-5 move. Such a graph corresponds to a triangulation of a 3-ball with five 4-simplices, and boundary a 3-sphere triangulated with five tetrahedra. The divergence of this graph can be associated to the vertex renormalization. In this case there are five vertices and the external data are ten spins and five tetrahedra. Let now $a, b, c = 1, \dots, 5$. To each couple $a < b$ is associated an internal intertwiner (tetrahedron); to each triple $a < b < c$ is associated an internal face (triangle shared by the three tetrahedra (ab) , (bc) and (ac)). Putting the external spins and intertwiners to zero, the spinfoam amplitude is

$$W_5 = \lambda^5 \sum_{j_{abc} i_{ab}} \prod_{a < b < c} d(j_{abc}) A_v(0, j_{1ab}, i_{1a}, 0) \dots A_v(0, j_{a5}, i_{a5}, 0), \quad (30)$$

where we have momentarily reinserted λ for clarity. As before, it reduces to an expression with $6j$ and $9j$ -symbols, and the intertwiners are evaluated at fixed values:

$$W_5 = \sum_{j_{abc}} \prod_{a < b < c} d(j_{abc}) 6j\left(\frac{1+\gamma}{2} j_{1ab}\right) 6j\left(\frac{1-\gamma}{2} j_{1ab}\right) \dots 6j\left(\frac{1+\gamma}{2} j_{a5}\right) 6j\left(\frac{1-\gamma}{2} j_{a5}\right) \prod_a f_{1a} \dots \prod_a f_{a5}. \quad (31)$$

Thus

$$W_5 \sim \sum_{j_{abc}} j^{10k} [(j^{-3/2})^2 (j^{-3/4})^4]^5, \quad (32)$$

and the degree of divergence is

$$D_{W_5} = 10 + 10k - 30. \quad (33)$$

The amplitude (31) converges for $k = 1$ and diverges logarithmically for $k = 2$. With the $\tilde{f}$ fusion coefficients

$$\widetilde{W}_5 \sim \sum_{j_{abc}} j^{10k} [(j^{-3/2})^2 (j^{-9/4})^4]^5 \quad (34)$$

and the degree of divergence is

$$D_{\widetilde{W}_5} = 10 + 10k - 60. \quad (35)$$

This converges for $k < 5$.

V. RESULTS

We have computed the naive degree of divergence of the first two interesting diagrams in loop quantum gravity, self-energy and vertex, for different choices of normalizations and face amplitude. With the fusion coefficients f defined in (7), the self-energy converges (naively) for $k < 1$ and the vertex converges (naively) for $k < 2$. Therefore with the $SO(4)$ measure factor (4) the self-energy diverges as a power and the vertex diverges logarithmically, similar to what happens in QED. With the $SU(2)$ measure factor (5) the self-energy diverges logarithmically while the vertex is finite. With the fusion coefficients $\tilde{f}$ defined in (8), there are no divergences for $k < 3$, therefore the two terms are finite with either of the two face amplitudes considered. These results are summarized in Table 1.

Fusion coefficients	Face amplitude	Self-energy	Vertex
f	$SU(2)(k = 1)$	logarithmic	finite
f	$SO(4)(k = 2)$	Λ^6	logarithmic
$\tilde{f}$	$SU(2)(k = 1)$	finite	finite
$\tilde{f}$	$SO(4)(k = 2)$	finite	finite

TABLE I: Naive degree of divergence (Λ is an infrared cut off).

The degree of divergence depends strongly on the normalization of face and vertex amplitudes. It is thus crucial to have better control over these quantities, which are not uniquely fixed in the literature.

All these divergences are in the infrared. The conventional ultraviolet divergences of gravity [19] do not appear in the theory; they are cut-off by the discretisation of the geometry given by the discrete character of the j_f spins. The divergences that appear have a clear geometrical interpretation: they correspond to “spikes” in the Regge triangulation, that is, very high “pyramids” that can spike out from a spacetime region bounded by a finite sphere.

We have not addressed the problem of characterizing *all* divergent diagrams. In particular: if the ball and the bubble diverge, would a regularized bubble make the ball finite as well? How many are the diagrams that one needs to regularize to make *all* of them finite? These issues and the implications of these results for loop quantum gravity, will be discussed elsewhere.

Acknowledgments

We are grateful to Elena Magliaro, Antonino Marcianò, Roberto Pereira, Daniele Pranzetti, Matteo Smerlak and Artem Starodubstev for numerous useful inputs and discussions.

-
- [1] J. Engle, R. Pereira, C. Rovelli: “The loop-quantum-gravity vertex-amplitude”, Phys. Rev. Lett., 99 (2007) 161301; arXiv:0705.2388.
 - [2] J. Engle, R. Pereira, C. Rovelli: “Flipped spinfoam vertex and loop gravity”, Nucl. Phys. B798 (2008) 251-290; arXiv:0708.1236.
R. Pereira, “Lorentzian LQG vertex amplitude”, Class. Quant. Grav. 25 (2008) 085013; arXiv:0710.5043.
 - [3] L. Freidel, K. Krasnov, “A New Spin Foam Model for 4d Gravity”. Class. Quant. Grav. 25 (2008) 125018, 2008; arXiv:0708.1595.
 - [4] E.R. Livine, S. Speziale, “A New spinfoam vertex for quantum gravity”, Phys. Rev. D76 (2007) 084028; arXiv:0705.0674.
E.R. Livine, S. Speziale, “Consistently Solving the Simplicity Constraints for Spinfoam Quantum Gravity,” Europhys. Lett. 81 (2008) 50004; arXiv:0708.1915.
 - [5] J. Engle, E.R. Livine, R. Pereira, C. Rovelli: “LQG vertex with finite Immirzi parameter” Nucl. Phys. B799 (2008) 136-149; arXiv:0711.0146.
 - [6] C. Rovelli, *Quantum Gravity* (Cambridge University Press, Cambridge 2004).
T. Thiemann, *Modern canonical quantum general relativity*, (Cambridge University Press, Cambridge, 2007).
 - [7] C. Rovelli, L. Smolin, “Knot theory and quantum gravity” Phys. Rev. Lett 61 (1988) 1155-1158.
C. Rovelli, L. Smolin, “Loop space representation for quantum general relativity”, Nucl. Phys. B331 (1990) 80-152.
A. Ashtekar, C. Rovelli, L. Smolin, “Weaving a classical geometry with quantum threads”, Phys. Rev. Lett 69 (1992) 237.
C. Rovelli, “A generally covariant quantum-field theory and a prediction on quantum measurements of geometry”, Nucl. Phys. B405 (1993) 797-815.
C. Rovelli, L. Smolin, “Discreteness of Area and Volume in Quantum Gravity”, Nucl. Phys. B442 (1995) 593-619; Nucl. Phys. B456 (1995) 734.

- [8] K. Wilson, “Confinement of Quarks”, Phys. Rev. D10 (1974) 2445.
- [9] C. Misner, “Feynman quantization of general relativity”, Rev. Mod. Phys. 29 (1957) 497.
S. W. Hawking, “The Path-Integral Approach to Quantum Gravity”, in *General Relativity: An Einstein Centenary Survey*, S. W. Hawking and W. Israel eds (Cambridge University Press, Cambridge 1979).
- [10] T. Regge, “General Relativity Without Coordinates”, Nuovo Cimento 19 (1961) 558-571.
- [11] A. Perez, “Finiteness of a spinfoam model for Euclidean quantum general relativity,” Nucl. Phys. B599 (2001) 427.
- [12] A. Perez, C. Rovelli: “A spin foam model without bubble divergences”, Nucl. Phys. B599 (2001) 255-282; gr-qc/0006107.
- [13] L. Crane, A. Perez, C. Rovelli: “Finiteness in spinfoam quantum gravity”, Phys. Rev. Lett. 87 (2001) 181301.
- [14] L. Crane, A. Perez, C. Rovelli: “A finiteness proof for the Lorentzian state sum spinfoam model for quantum general relativity”, gr-qc/0101088.
- [15] E. Magliaro, C. Perini, C. Rovelli, “Numerical indications on the semiclassical limit of the flipped vertex, Class. Quant. Grav. 25 (2008) 095009, arXiv:0710.5034.
B. Dittrich, S. Speziale, “Area-angle variables for general relativity,” New J. Phys. 10 (2008) 083006; arXiv:0802.0864.
E. Alesci, E. Bianchi, E. Magliaro, C. Perini, “Intertwiner dynamics in the flipped vertex, arXiv:0808.1971.
E. Bianchi, A. Satz, “Semiclassical regime of Regge calculus and spin foams, arXiv:0808.1107.
F. Conrady, L. Freidel, “On the semiclassical limit of 4d spin foam models, arXiv:0809.2280.
- [16] M.P. Reisenberger, C. Rovelli, “Spacetime as a Feynman diagram: The connection formulation,” Class. Quant. Grav. 18 (2001) 121.
- [17] D. Oriti, “The group field theory approach to quantum gravity,” in *Approaches to Quantum Gravity - toward a new understanding of space, time, and matter*, D. Oriti editor (Cambridge University Press 2008). arXiv:gr-qc/0607032.
L. Freidel, “Group Field Theory: an overview”, Int. Journ. Theor. Phys. 44 (2005) 1769-1783.
R. DePietri, L. Freidel, K. Krasnov, C. Rovelli “Barrett-Crane model from a Boulatov-Ooguri field theory over a homogeneous space”, Nucl. Phys. B574 (2000) 785-806.
- [18] G. Ponzano, T. Regge, “Semiclassical limit of Racah coefficients. in *Spectroscopic and Group Theoretical Methods in Physics* F. Block editor (North Holland, Amsterdam, 1968).
- [19] S. Weinberg, “Ultraviolet divergences in quantum theories of gravitation”, in *General Relativity: An Einstein Centenary Survey*, S. W. Hawking and W. Israel eds (Cambridge University Press, Cambridge 1979).
- [20] E. Alesci, E. Bianchi, E. Magliaro, C. Perini, “Asymptotics of LQG fusion coefficients”, arXiv:0809.3718.
- [21] I. Khavkine, “Evaluation of new spin foam vertex amplitudes, arXiv:0809.3190.
- [22] A. Yutsin, I. Levinson, V. Vanagas, *Mathematical Apparatus of the Theory of Angular Momentum* (Israel Program for Scientific Translation, Jerusalem, 1962).
D. Varshalovich, A.N. Moskalev, and V.K. Khersonskii, *Quantum Theory of Angular Momentum* (World Scientific, Singapore 1988).

APPENDIX A: NORMALIZATIONS

The Wigner 3j symbols

$$i^{m_1 m_2 m_3} \equiv \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \quad (\text{A1})$$

are real trivalent intertwiners, namely invariant tensors in the tensor product $H_{j_1} \otimes H_{j_2} \otimes H_{j_3}$ of three $SU(2)$ representations, normalized by

$$\sum_{m_1 m_2 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = 1. \quad (\text{A2})$$

They are related to the Clebsch-Gordan coefficients by a normalization factor

$$\begin{pmatrix} j_1 & j_2 & J \\ m_1 & m_2 & -M \end{pmatrix} = \frac{(-1)^{j_1 - j_2 - M}}{\sqrt{2J + 1}} \langle j_1 j_2, m_1 m_2 | JM \rangle. \quad (\text{A3})$$

The Wigner 3j symbols as usually represented by the vertex of three lines that join.

$$i^{abc} = \begin{array}{c} b \\ | \\ \bullet \\ / \quad \backslash \\ a \quad c \end{array} . \quad (\text{A4})$$

Joining free ends of these lines indicates contraction with the intertwiner g_{ab} between a representation and its dual. The 4-valent intertwiners can be written in terms of trivalent intertwiners. An orthonormal basis of 4-valent intertwiners is given by

$$i_J^{abcd} \equiv \sqrt{2J+1} \sum_{mn} g_{mn} \begin{pmatrix} j_1 & j_2 & J \\ a & b & m \end{pmatrix} \begin{pmatrix} J & j_3 & j_4 \\ n & c & d \end{pmatrix} \equiv \sqrt{2J+1} i^{abm} i_m^{cd}. \quad (\text{A5})$$

Orthonormality can be proven explicitly by writing

$$\begin{aligned} i_J^{abcd} i_{Kabcd} &= \sqrt{2J+1} \sqrt{2K+1} g_{aa'} g_{bb'} g_{cc'} g_{dd'} g_{mn} g_{m'n'} \begin{pmatrix} j_1 & j_2 & J \\ a & b & m \end{pmatrix} \begin{pmatrix} J & j_3 & j_4 \\ n & c & d \end{pmatrix} \begin{pmatrix} j_1 & j_2 & K \\ a' & b' & m' \end{pmatrix} \begin{pmatrix} K & j_3 & j_4 \\ n' & c' & d' \end{pmatrix} \\ &= \sqrt{2J+1} \sqrt{2K+1} \delta_{JK} \frac{\delta_n^m \delta_m^n}{(2J+1)(2K+1)} = \delta_{JK}. \end{aligned} \quad (\text{A6})$$

The standard graphical notation for the 4-valent intertwiners indicates $i^{abm} i_m^{cd}$, and not the normalized intertwiners $i_J^{abcd} = \sqrt{2J+1} i^{abm} i_m^{cd}$, that is

$$i_J^{abcd} = \sqrt{2j+1} \begin{array}{c} b \quad \quad c \\ \quad \diagdown \quad \diagup \\ \quad \bullet \quad \bullet \\ \quad \diagup \quad \diagdown \\ a \quad \quad d \end{array} . \quad (\text{A7})$$

Since the last form an orthonormal basis on the invariant part of the product of four representation spaces, we have

$$\sum_J i_J^{abcd} i_J^{a'b'c'd'} = \sum_J i_J^{acbd} i_J^{a'c'b'd'} \quad (\text{A8})$$

from which the recoupling theorem follows easily. In terms of these quantities, and the unitary representation matrices of $\text{SU}(2)$, $R^{(j)a}_b(U)$, we have

$$\int dU R^{(j)a}_b(U) R^{(k)c}_d(U) = \frac{1}{2j+1} \delta^{jk} g^{ac} g_{bd} \quad (\text{A9})$$

$$\int dU R^{(j)a}_d(U) R^{(k)b}_e(U) R^{(l)c}_f(U) = i^{abc} i_{def} \quad (\text{A10})$$

$$\int dU R^{(j)a}_e(U) R^{(k)b}_f(U) R^{(l)c}_g(U) R^{(m)d}_h(U) = \sum_J i_J^{abcd} i_{Jefgh} \quad (\text{A11})$$

and

$$\delta(U) = \sum_j (2j+1) \text{tr}[R^{(j)}(U)] \quad (\text{A13})$$

which are the basic formulas for deriving the spinfoam representation of BF theory. Expanding the BF partition function

$$Z = \int dU_l \prod_f \delta(U_{f_1} \dots U_{f_n}) \quad (\text{A14})$$

in representations and using the formulas above we obtain the well known Ponzano-Regge expression

$$Z = \sum_{j_f} \prod_f (2j_f+1) \prod_v 6j(j_f) \quad (\text{A15})$$

in the 3d case. While in the (SU(2)) 4d case we obtain

$$Z = \sum_{j_f, i_e} \prod_f (2j_f + 1) \prod_v A(j_f, i_e), \quad (\text{A16})$$

where the vertex amplitude is the contraction of five *normalized* intertwiners.

$$A(j_f, i_e) = \otimes_a i_a. \quad (\text{A17})$$

This is not the 15j symbol as usually defined in representation theory texts:

$$15j(j_{ab}, j_a) \equiv \begin{Bmatrix} j_{12} & j_{23} & j_{34} & j_{45} & j_{51} \\ j_{35} & j_{41} & j_{52} & j_{13} & j_{25} \\ j_1 & j_2 & j_3 & j_4 & j_5 \end{Bmatrix} \quad (\text{A18})$$

but rather

$$A(j_{ab}, j_a) = 15j_N(j_{ab}, j_a) \equiv \sqrt{\prod_a (2j_a + 1)} 15j(j_{ab}, j_a). \quad (\text{A19})$$

APPENDIX B: ASYMPTOTICS OF FUSION COEFFICIENTS WITH A SINGLE ZERO SPIN

When one of the four spins is zero, say j_4 , the fusion coefficients reduce, up to a sign, to

$$\tilde{f}_{i^+ i^-}^i(j_1, j_2, j_3, 0) = \pm \delta_{i^+, j_3^+} \delta_{i^-, j_3^-} \delta_{i, j_3} 9j(j_1, j_2, j_3), \quad (\text{B1})$$

where

$$9j(j_1, j_2, j_3) \equiv \begin{Bmatrix} j_1^+ & j_3^+ & j_2^+ \\ j_1^- & j_3^- & j_2^- \\ j_1 & j_3 & j_{j_2} \end{Bmatrix} \quad (\text{B2})$$

is a Wigner $9j$ symbol. We are interested in the asymptotic behavior of (B1) for large spins. (For a numerical approach, see [21].) The $9j$ -symbol with two degenerate columns can be written as [22]

$$\begin{Bmatrix} a & f & c \\ b & g & d \\ a+b & h & c+d \end{Bmatrix} = (-1)^{f-g+a+b-(c+d)} \begin{pmatrix} f & g & h \\ a-c & b-d & -(a+b-(c+d)) \end{pmatrix} \times \quad (\text{B3})$$

$$\times \sqrt{\frac{(2a)!(2b)!(2c)!(2d)!(a+b+c+d-h)!(a+b+c+d+h+1)!}{(2a+2b+1)!(2c+2d+1)!(a+c-f)!(a+c+f+1)!(b+d-g)!(b+d+g+1)!}}. \quad (\text{B4})$$

The degenerate $3j$ -symbol, in which the third spin is the sum of the first two can be derived easily from well-known expressions for the $3j$ symbol [22]:

$$\begin{pmatrix} f & g & f+g \\ m & n & -(m+n) \end{pmatrix} = (-1)^{f-g+m+n} \sqrt{\frac{(2f)!(2g)!(f+g-m-n)!(f+g+m+n)!}{(1+2f+2g)!(f-m)!(f+m)!(g-n)!(g+n)!}}. \quad (\text{B5})$$

We can use the representations (B3) and (B5) both for $0 \leq \gamma < 1$ and for $\gamma > 1$. This is because in the first case we have $j^+ + j^- = j$; while in the second have $j + j^- = j^+$: in either case there is one entry which is given by the sum of the others. Both in (B3) and (B5) some factorials are written in the form $(1+x)!$; we substitute them with the expression $(1+x)x!$. Then, as we are interested in the large spin behavior, we apply Stirling's approximation

$$x! \sim \sqrt{2\pi x} e^{-x+x \log x}$$

to all factorials in the resulting expression. All factors coming from the exponential in the Stirling formula cancel out, giving 1. The only contribution is provided by the monomials $(1+x)$ we factored out, and from $\sqrt{2\pi x}$ in Stirling's expression. So we obtain easily the asymptotic expression

$$9j(j_1, j_2, j_3) \sim \pm \beta_\gamma \frac{\sqrt{2}}{\pi^{1/4}} \left[\frac{1}{\sqrt{j_1 j_2 j_3 (j_1 + j_2 - j_3)(j_3 + j_1 - j_2)(j_3 + j_2 - j_1)(j_1 + j_2 + j_3)^3}} \right]^{1/2} \quad (\text{B6})$$

where

$$\beta_\gamma = \begin{cases} \left(\frac{1}{1-\gamma^2}\right)^{1/4} & 0 \leq \gamma < 1 \\ \left(\frac{1}{\gamma^2-1}\right)^{3/4} & \gamma > 1 \end{cases} \quad (\text{B7})$$

Therefore the $9j$ Wigner symbols that appear in the fusion coefficients behave like $j^{-9/4}$ for large spins. Accordingly the fusion coefficients behave as

$$f \sim \frac{1}{j^{3/4}}, \quad \tilde{f} \sim \frac{1}{j^{9/4}}. \quad (\text{B8})$$

These are the asymptotic expressions used in the text.